

# ABDOMINAL PAIN STANDING ORDER

- Initiate Immediate Supportive Care
- O2 to keep sat >90%
  - Cardiac Monitor (if available)

I  
N  
C  
L

Use standing order on patients greater than 16 years of age complaining of abdominal pain

E  
X  
C  
L  
U  
S  
I  
O  
N

- Do not implement standing order on patients with the following symptoms:
- Pregnancy
  - Meeting trauma triage criteria
  - Abdominal trauma
  - Mixed symptoms
  - Known history of cardiac disease

## STABLE

SBP >90 or HR <110

- Assess acuity and severity of pain
- Transport in position of comfort with supportive measures as indicated

If patient's condition deteriorates notify receiving facility of change in status. Consider transport to closest facility

MEDS Notification to receiving facility:  
Advise if patient is *Stable* or *Unstable*

## UNSTABLE

SBP <90 or HR >110

- Follow STABLE patient orders
- Initiate a large bore IV of Normal Saline
- If the patient is pale/diaphoretic, or has a systolic SBP less than 90 or a heart rate greater than 110 administer 20 ml/kg Normal Saline
- Reassess hemodynamic and pulmonary status and rebolus as indicated
- For nausea or vomiting administer Ondansetron:  
**ADULT: IV 4mg over 2-5 min. May repeat once after 15 min if no response.**

Consider medical direction contact for analgesic orders.