

ALLERGIC REACTION

Initiate Immediate Supportive Care:

- Keep O2 sat > 90%
- Cardiac Monitor

I
N
C
L
U
S
I
O
N

Stable Allergic Reaction

- Urticaria (Hives)
- Sense of Dyspnea
- Sense of Oropharyngeal Swelling
- Sense of Throat Tightness

I
N
C
L
U
S
I
O
N

Unstable Allergic Reaction

- Signs of Shock
- Objective Signs of **severe** Respiratory Distress
- Objective signs of Airway Compromise

O
R
D
E
R
S

STABLE ALLERGIC REACTION

- Establish IV NS TKO
- Administer Diphenhydramine IVP 1mg/kg up to 25 mg

If patient has mixed symptoms or wishes to refuse contact medical direction.

MEDS Notification to receiving facility: Advise if patient is Stable or Unstable

UNSTABLE ALLERGIC REACTION

- Manage Airway
- Administer Epinephrine:

BLS	ALS
<ul style="list-style-type: none"> • Administer Epi pen (pediatric auto-injector if \leq 30 kg) 	<ul style="list-style-type: none"> • 1:10,000 IV/IO 0.5 mg repeat every 2 -5 minutes for patients in extremis OR • 1:1,000 IM (ant thigh preferred) 0.01mg/kg up to 0.5 mg total \pm

O
R
D
E
R
S

- If wheezing, give Albuterol one unit dose with Ipratropium one unit dose via SVN
- Initiate IV NS to keep SBP > 90
- Administer Diphenhydramine IVP 1mg/kg up to 25 mg total
- Administer Methylprednisolone 2 mg/kg IVP up to 125 mg
- If Methylprednisolone unavailable administer Dexamethasone 0.5mg/kg IVP up to 20mg