

ALLERGIC REACTION STANDING ORDER

Initiate Immediate Supportive Care

- Keep O2 sat > 90%
- Cardiac Monitor

Inclusion: Stable Allergic Reaction

- Urticaria (Hives)
- Sense of Dyspnea
- Sense of Oropharyngeal Swelling
- Sense of Throat Tightness

Inclusion: Unstable Severe Allergic Reaction

- Signs of Shock
- Objective Signs of Respiratory Distress: Wheezing or Stridor
- Objective signs of Airway Compromise

EXCL: Patient wishes to refuse

Contact medical direction, **IMMEDIATELY**

EXCL: Patient has Mixed Symptoms or wishes to refuse

Orders: STABLE ALLERGIC REACTION

(Check Medication Allergies)

- Establish IV NS TKO
- Administer Benadryl IVP 1mg/kg up to 25 mg

Orders: UNSTABLE ALLERGIC REACTION

(Check Medication Allergies)

- Manage Airway
- Administer Epinephrine
 - Administer Epi pen (pediatric auto-injector if ≤ 30 kg)
 - 1:1,000 IM (ant thigh preferred) 0.01ml/kg up to 0.5 ml total*
- If wheezing, give Albuterol one unit dose with Atrovent one unit dose via SVN
- Initiate IV NS to keep SBP > 90
- Administer Benadryl IVP 1mg/kg up to 25mg total
- Administer Solumedrol 2mg/kg IVP up to 125 mg total
- If Solumedrol unavailable, administer Decadron 0.5mg/kg IVP up to 20mg

*Use caution in patients with known Ischemic Cardiac Disease

MEDS notification to receiving facility: Advise if patient is Stable or Unstable